arXiv:1212.5933v1 [quant-ph] 24 Dec 2012

State (in)dependent graphs: fractional colorings and a cloud of contextuality

Ravishankar Ramanathan[1, 2, 3, *] and Pawel Horodecki[2, 4]
[1] University of Gdansk 80-952, Gdansk, Poland
[2] National Quantum Information Center of Gdansk, 81-824 Sopot, Poland
[3] Centre for Quantum Technologies, National University of Singapore, 3 Science Drive 2, 117543 Singapore, Singapore
[4] Faculty of Applied Physics and Mathematics, Technical University of Gdansk, 80-233 Gdansk, Poland

Contextuality of non-perfect orthogonality graphs is studied and shown to occur even in cases when the Lovasz and independence numbers are equal. It is proven that for given dimension (i) contextuality of a quantum state is entirely determined by its spectrum, hence pure and maximally mixed states represent the two extremes of contextual behavior and (ii) state-independent contextuality (S-IC) is equivalent to contextuality of the maximally mixed state. A necessary and sufficient condition for a graph to be S-IC is provided in terms of the fractional chromatic number $\chi_f(G)$ and is shown to be stronger than its direct analog in terms of the (standard) chromatic number $\chi(G)$. The completion paradigm is then studied in which extra projections required to physically realize a complete measurement are shown to constitute a "cloud of contextuality" which may render the full set of projectors S-IC even if the original set of projectors was not. On that new level, the intriguing interplay between the $\chi_f(G)$ and $\chi(G)$ conditions is shown to lead to a striking property of the well-known Kneser graphs and subgraphs in regard to their quantum realization.

Introduction. One of the most striking features of the quantum world is contextuality, the notion that outcomes cannot be assigned to measurements independent of the particular contexts in which the measurements are realized. The incompatibility of such assignments manifests itself in the famous Kochen-Specker paradox [1], for every quantum system belonging to a Hilbert space of dimension greater than two, irrespective of its actual state, a finite set of measurements exists whose results cannot be assigned in a context-independent manner. This phenomenon is known as state-independent contextuality, there are also state-dependent tests [2] in which only a subset of quantum states in a certain Hilbert space displays contextual behavior while the results of measurements on other states can be explained by non-contextual assignments (or their probabilistic mixtures).

A natural question is: when does a given set of (projective) measurements reveal the contextuality of some quantum state i.e. when is a set of measurements contextual? Moreover, one may also refine the question to: when does a set of (projective) measurements reveal the contextuality of all quantum states (belonging to a particular Hilbert space of dimension d)? A lot of attention has been devoted to these questions [3, 4], in particular to the problem of finding the minimal set of measurements that reveals the contextuality of all states of a given dimension [5–7]. It has been shown that the originally studied K-S sets are not the only class of state-independent contextual measurements [8], so that the identification of general conditions has gained importance, more so with the application of contextuality to device-independent security [9].

In the study of contextual measurements, a number of tools and invariants of graph theory have appeared (see [10] for a recent approach). In this paper, we focus on orthogonality graphs where each projective measurement is assigned a vertex of the graph and edges connect commuting (orthogonal) measurements. Regarding state-depending contextuality, a certain condition for a graph to be non-contextual in terms of its Lovasz theta number and independence number is shown to not always hold by means of a counter-example. The state-dependent contextuality of a proper subgraph is shown to guarantee the contextuality of the entire graph so that an investigation of the class of odd cycle graphs and their complements provides much information about the class of all contextual graphs. The quantitative measure of contextuality for any quantum state [11] is shown to depend only on the spectrum of its eigenvalues so that among all states belonging to a given dimension, the maximally mixed state is the least and the pure states are the most contextual. This fact is used to formulate a necessary and sufficient condition in terms of the fractional chromatic number for an orthogonality graph to reveal state-independent contextuality. Additional projectors required to physically realize complete measurements are shown to be capable of forming a "cloud of contextuality" that reveals state-independent contextuality even when the original set of projectors is unable to do so. Proofs of propositions are collected in the Supplementary Material.

The class of contextual graphs. We focus on orthogonality graphs for a given set of projective measurements $\{\Pi_k\}$, commonly the projectors are of rank one ($\Pi_k = |v_k\rangle\langle v_k|$). In the orthogonality graphs, every projective measurement is denoted by exactly one vertex in the graph and edges connect two vertices if and only if the corresponding projectors are orthogonal. A classical (non-contextual) assignment of outcomes $\{0, 1\}$ is then an assignment of 1's to the vertices belonging to an independent set of the graph (a set of mutually disconnected vertices) and 0's otherwise. This assignment guarantees that in each orthonormal basis only one projector is assigned the value 1. A set of projectors is said to realize graph G if it strictly obeys the orthogonality constraints of G. Given a realization of G, it is natural to allow a global unitary transformation on all the projectors since these preserve G. We therefore investigate the contextual properties of orthogonality graphs by considering an optimization over all projectors that realize the particular graph. Note that other representations of contextual measurements by graphs can be translated to orthogonality graphs. For instance, a reformulation of the representation via hypergraphs [6, 11] with edges consisting of commuting observables is to interpret the common eigenstates of every commuting set of observables as projectors realizing the orthogonality graph.

For an orthogonality graph G with vertex set $\mathcal{V} = \{v_1, \dots v_n\}$, let $\mathcal{I}(G)$ denote the set of all independent sets $I^{(j)}(G)$ of G. One may construct corresponding (incidence) vectors $\vec{I}^{(j)}(G) \in \mathtt{R}^n$, $j \in \{1, \dots, |\mathcal{I}(G)|\}$ with components $\vec{I}^{(j)}_k(G) = 1$ if $v_k \in I^{(j)}(G)$ and $\vec{I}^{(j)}_k(G) = 0$ otherwise. The convex hull of all such vectors $\vec{I}^{(j)}(G)$ then corresponds to the classical non-contextual polytope for the graph G (also known as the stable-set polytope STAB(G)) [12]. The set of probabilities realizable in general probabilistic theories consistent with the no-disturbance principle [13] (in which probabilities are well-defined and context independent) reside in the polytope known as the clique-constrained stable set polytope QSTAB(G). The notion of clique-constraint refers to the fact that the sum of probabilities for projectors in every orthonormal basis must not exceed unity (a clique refers to a complete subgraph in which all vertices are connected and represents a basis). The set of quantum probabilities is in general not a polytope but is a subset of QSTAB(G). A quantum state ρ is represented by a vector $\vec{x}^{\{\rho\}}(G) \in \mathtt{R}^n$ with components $\vec{x}^{\{\rho\}}_k(G) = Tr[\Pi_k \rho]$.

Definition 1. An orthogonality graph G is said to be contextual for dimension d if $\exists \rho \in \mathcal{H}^d$ such that $\vec{x}^{\{\rho\}}(G) \notin STAB(G)$. A graph that is not contextual for any dimension is called non-contextual. G is said to be state-independent contextual (S-IC) for dimension d if $\forall \rho \in \mathcal{H}^d$, $\vec{x}^{\{\rho\}}(G) \notin STAB(G)$.

It is well-known that the class of perfect graphs has STAB(G) = QSTAB(G) [14] so that such graphs cannot be contextual. They are defined by the notion that the chromatic number of any induced subgraph is equal to its clique number, $\chi(H) = \omega(H)$ for $H \subseteq G$. Chromatic number $\chi(G)$ is the number of colors needed to properly color the vertices of the graph with adjacent vertices being assigned different colors and clique number $\omega(G)$ is the size of its largest clique. By the Strong Perfect Graph Theorem [15], perfect graphs are precisely those that do not contain as induced subgraphs any odd cycle of five or more vertices or its complement (the complement $\bar{G}$ of G has the same vertex set as G, two vertices are connected by an edge in $\bar{G}$ iff they are not in G). It follows that all contextual orthogonal graphs contain induced odd cycles (C_{2k+1} with $k \geq 2$) or their complements $\bar{C}_{2k+1}$ [3].

The question then arises: does the presence of C_{2k+1} or $\bar{C}_{2k+1}$ in G guarantee that G is contextual, i.e. are all non-perfect graphs contextual? One possible condition [3] for an orthogonal graph to be non-contextual is that its Lovasz theta number $\theta(G)$ equal its independence number $\alpha(G)$. The Lovasz theta number $\theta(G)$ [16] of graph G on n vertices is defined as $\theta(G) := \max_{\{\Pi_i\}} \| \sum_{i=1}^{n} \Pi_i \|_\infty$ where the maximization is over all sets of projectors $\{\Pi_i\}$ realizing the graph and infinity norm is the maximal eigenvalue. The independence number $\alpha(G)$ is the size of the maximal independent set in G. The intuition behind this condition [12] is that it excludes the violation by any quantum state of known inequalities of the kind $\sum_{i=1}^{n} \langle \Pi_i \rangle \leq \alpha(G)$, where the braces denote expectation values. The fact that there exist non-perfect graphs for which $\theta(G) = \alpha(G)$ would then be evidence that the class of perfect graphs is strictly smaller than the class of non-contextual graphs.

However, it can be shown that this condition does not pick out the class of all non-contextual graphs according to the definition above, if we restrict ourselves as here to simple, undirected and non-weighted graphs. For instance, there are non-perfect graphs (see Supplementary Material) for which $\theta(G) = \alpha(G)$ that can be shown to be contextual. Therefore, while $\alpha(G) < \theta(G)$ guarantees a contextual graph, $\alpha(G) = \theta(G)$ does not imply non-contextuality. One step in identifying the class of contextual graphs is the following observation.

Observation 1. If in any realization of orthogonal graph G, some proper subgraph H of G is contextual, then G is contextual.

Since the odd cycles C_{2k+1} or their complements $\bar{C}_{2k+1}$ (with integral $k \geq 2$) are always present as induced subgraphs in non-perfect graphs, one may ask if in any realization these graphs are contextual.

Observation 2. Non-perfect graphs containing induced C_5 are contextual in dimension 3.

The question whether all non-perfect graphs can be contextual merits further investigation. Here, we proceed with identifying the subclass of state-independent graphs.

Measure of contextuality and spectra of states. Given an orthogonality graph G realizable by a set of projectors $\{\Pi_j\}$ with $\Pi_j \in \mathcal{H}^d$, a quantitative measure [11] enables us to study the relative contextuality of quantum states $\rho \in \mathcal{H}^d$. Here, we show that the measure of contextuality of a quantum state only depends on its spectrum. As a consequence, for any G we show that the maximally mixed state $\mathbb{1}_d$ is the least contextual while the pure states display maximum contextuality.

Allowing a maximization over all sets of projectors $\{\Pi_j\}$ realizing G, for given ρ the distance of $\vec{x}^{\{\rho\}}(G)$ from STAB(G) defines a measure of contextuality $M^{(G)}(\rho)$

given by

$$\begin{aligned} M^{(G)}(\rho) &:= \max_{\{\Pi_k\}} M^{(G)}_{\{\Pi_k\}}(\rho) \qquad (1)\\ &:= \max_{\{\Pi_k\}} \min_{p(\lambda)} \sum_{c\in\mathcal{C}(G)} \frac{1}{|\mathcal{C}(G)|} D(q^{(\rho,\{\Pi_k\})}(\lambda_c)||p(\lambda_c)). \end{aligned}$$

Here, the maximization is over all sets of projectors realizing G, $\mathcal{C}(G)$ denotes the set of maximal cliques in the graph G and $q^{(\rho,\{\Pi_k\})}(\lambda_c) = \{Tr[\rho\Pi_k^{(c)}]\}$ is the probability distribution corresponding to the clique (context) c. The minimization is performed over the (classical) joint probability distributions $p(\lambda)$ for all contexts in the graph G with $p(\lambda_c)$ being the corresponding marginal distribution for context c. The relative entropy distance is $D(q(\lambda_c)||p(\lambda_c)) = \sum_i q(\lambda_c)_i \log \frac{q(\lambda_c)_i}{p(\lambda_c)_i}$ where summation extends over all projectors belonging to the context c and any (possibly different rank) projectors needed to realize a complete measurement ($\sum_i q(\lambda_c)_i = 1$). Let $\vec{\kappa}(\rho)$ denote the spectrum of ρ, we then have the following.

Theorem 1. $M^{(G)}(\rho)$ is entirely determined by $\vec{\kappa}(\rho)$. Moreover, $\forall \rho_1, \rho_2 \in \mathcal{H}^d$ and $\forall G$, $\vec{\kappa}(\rho_1) \prec \vec{\kappa}(\rho_2) \Rightarrow M^{(G)}(\rho_1) \leq M^{(G)}(\rho_2)$. In particular, $M^{(G)}(\mathbb{1}_d) \leq M^{(G)}(\rho)$ $\forall \rho \in \mathcal{H}^d$.

Here $\vec{\kappa}(\rho_1) \prec \vec{\kappa}(\rho_2)$ denotes that the vector of eigenvalues of ρ_2 majorizes that of ρ_1. Note that the violation of non-contextual inequalities (defining the facets of STAB(G)) by quantum states also follows the majorization order, i.e., $\vec{\kappa}(\rho_1) \prec \vec{\kappa}(\rho_2)$ implies that the violation of any non-contextual inequality by ρ_1 does not exceed that by ρ_2. Knowing that the maximally mixed state is the least contextual among all states in the given Hilbert space, the identification of state-independent measurements can be reduced to the question: for given G, is $\vec{x}^{\{\mathbb{1}_d\}(G)} \notin STAB(G)$? As we shall see, there is one graph-theoretic quantity (the fractional chromatic number) that precisely identifies when this holds.

State-independent contextual graphs. Since non-perfect graphs are the only contextual graphs, a preliminary consideration is whether for any $k \geq 2$, C_{2k+1} or $\bar{C}_{2k+1}$ are S-IC graphs, to this end we have.

Proposition 1. C_{2k+1} and $\bar{C}_{2k+1}$ are not S-IC graphs $\forall k \geq 2$, $k \in \mathcal{Z}$.

We now proceed to formulate a necessary and sufficient condition for an orthogonal graph G realizable in dimension d to be state-independent contextual in that dimension in terms of its fractional chromatic number $\chi_f(G)$ [19], defined below using the notion of fractional colorings. We show that this condition is stronger than an analogous condition proposed in [4] in terms of the chromatic number $\chi(G)$.

Definition 2.A fractional coloring of a graph G [19] is a non-negative real-valued function f on $\mathcal{I}(G)$ such that for any vertex v of G, $\sum_{s\in\mathcal{I}(G,v)} f(s) \geq 1$, where $\mathcal{I}(G,v)$ denotes the set of independent sets of G that contain vertex v. The weight of a fractional coloring is $w = \sum_{j=1}^{\mathcal{I}(G)} f(I^{(j)}(G))$, and $\chi_f(G) = \min w$. Equivalently, $\chi_f(G) = \min \frac{a}{b}$ s.t. there is a coloring of G using a colors that assigns b colors to each vertex with adjacent vertices getting disjoint sets of colors.

For any graph $\omega(G) \leq \chi_f(G) \leq \chi(G)$. The first inequality comes from the fact that in a clique of size ω assigning b colors to each vertex requires at least $\omega(G)b$ colors; the second because $\chi(G)$ is the particular restriction $b = 1$ to the defintion above. **Theorem 2.** An orthogonal graph G realizable by a set of rank-r projective measurements $\{\Pi_j\}$ with $\Pi_j \in \mathcal{H}^d$ is state-independent contextual for dimension d if and only if $\chi_f(G) > \frac{d}{r}$.

In [4], it was shown that a set of rank-1 projective measurements reveals the contextuality of the maximally mixed state if and only if $\chi(G) > d$. While this latter condition could hold in the"completion paradigm" (in next section) it is not sufficient in the standard approach focusing on the absence of $\vec{x}^{\{\mathbb{1}_d\}}(G)$ in STAB(G). One counter-example is provided by the graph formed by the join of two copies of the orthogonality graph G_{YO} considered in [8]. The graph G_{YO} is a 13-vertex graph with $\chi(G_{YO}) = 4$ and $\xi(G_{YO}) = 3$. Here $\xi(G)$ denote the smallest dimension d in which the graph G may be realized by rank-1 projectors with distinct vertices assigned distinct projectors and two vertices connected by an edge iff the corresponding projectors are orthogonal. The join of two graphs is their graph union with additional edges connecting all the vertices of one graph with those of the other. The join (J) of two G_{YO} graphs has $\chi(J(G_{YO}, G_{YO})) = 8$ and $\xi(J(G_{YO}, G_{YO})) = 6$. Moreover, its fractional chromatic number can be computed by means of a linear program to be $\chi_f(J(G_{YO}, G_{YO})) = 6\frac{4}{11}$ so that in dimension 7 this graph is not by itself S-IC as $\vec{x}^{\{\mathbb{1}_7\}}(J(G_{YO}, G_{YO})) \in STAB(J(G_{YO}, G_{YO}))$ (see Supplementary Material). Therefore, the condition $\chi(G) > d$ is not sufficient for a graph G to reveal state-independent contextuality by itself. Moreover, the gap between $\chi_f(G)$ and $\chi(G)$ can be arbitrarily large [19] so that $\chi_f(G)$ is by itself a better indicator of the S-IC property of G. In general, $\chi_f(G)$ is computed as a lower bound to $\chi(G)$ by means of the fractional relaxation to the integer linear program required to compute the latter quantity. This does not imply that $\chi_f(G)$ can be computed in polynomial time, since the number of independent sets may be exponential in the number of vertices. Since $\chi(G)$ upper bounds $\chi_f(G)$, $\chi(G) > d$ is only a necessary condition for G to be state-independent contextual. As we shall see in the next section, it may constitute a sufficient condition for a graph G_c, formed from G by adding projectors so that each clique constitutes a complete orthonormal basis, to be state-independent contextual, although this remains to be shown. Before proceeding to the "completion paradigm", let us derive a bound on the set of dimensions in which G can by itself be an S-IC graph.

Proposition 2. An orthogonal graph G can be state-independent contextual by itself only for dimensions $\xi(G) \leq d < 2^{\xi(G)-1}$.

Completion paradigm and a cloud of contextuality. We have seen that $\chi_f(G) > d$ is necessary and sufficient for $\vec{x}^{\{\mathbb{1}_d\}}(G) \notin STAB(G)$, therefore for graph G to be S-IC in dimension d. While usually considered orthogonal graphs do not have all cliques of the same size, we note that to physically realize the measurements, it is imperative that each context correspond to a complete measurement. In this paradigm, the orthogonal graph G must be completed by additional projectors to a new graph G_c in which each clique (context) is of the same size (equal to dimension d). Note that such a completion by rank-1 projectors is by no means unique,an optimization procedure is needed to complete G to an S-IC G_c. It may thus happen that G is by itself not an S-IC graph while some completion G_c is. We refer to this as a cloud of contextuality around the projectors in G.

Given the basis-completed graph G_c (in which all cliques are of the same size $\omega(G_c) = d$), $\chi_f(G_c) > d$ is necessary and sufficient for G_c to be S-IC. One may consider whether this condition is equivalent to $\chi(G_c) > d$, i.e. are there graphs G_c for which $\omega(G_c) = \chi_f(G_c) = d$ and $\chi(G_c) > d$? A particular candidate for this task is the well-known class of Kneser graphs $K(nb : b)$ with $n \geq 2$ and $b \geq 2$. These graphs are obtained by the following construction: consider an nb element set and obtain all b element subsets, each of these subsets corresponds to a single vertex of the graph and an edge connects two vertices iff they correspond to disjoint subsets. These graphs are basis-complete since all cliques are of the same size $\omega(K(nb : b)) = n$. Moreover, $\chi_f(K(nb : b)) = n$ and $\chi(K(nb : b)) = (n-2)b + 2$ [19] so that the separation between χ_f and χ can be made arbitrarily large by increasing b. Then $\chi_f(G_c) > d \Leftrightarrow \chi(G_c) > d$ would imply the following striking restriction on the quantum realization of these graphs, namely: Any basis-complete graph G_c for which $\chi_f(G_c) = \omega(G_c)$ and $\chi(G_c) > \omega(G_c)$ cannot be quantumly realized in dimension $\omega(G_c)$. Towards this surprising consequence, we prove the following.

Proposition 3. All basis-complete graphs G_c for which $\chi_f(G_c) = \omega(G_c)$ ($= n$, say) are proper subgraphs of the Kneser graphs $K(nb : b)$ for some integral b. $K(nb : b)$ with $n \geq 2$ and $b \geq 2$ themselves cannot be quantumly realized by rank-1 projectors in dimension n. Moreover, for $b = 2$, all proper basis-complete subgraphs of $K(2n : 2)$ also cannot be realized in dimension n.

While $\chi_f(G) > d \Rightarrow \chi_f(G_c) > d$ since the fractional chromatic number of G_c cannot be smaller than that of its subgraph G, it is not clear if $\chi(G) > d \Rightarrow \chi_f(G_c) > d$. One may conjecture based on Proposition 3 that the latter implication does indeed hold. Finally, note that neither $\chi_f(G) > d$ nor $\chi(G) > d$ are necessary for G_c to be S-IC. As an example, consider the graph $G_{YO} \setminus \{1\}$ formed by deleting one vertex from the state-independent graph in Fig. (2). $G_{YO} \setminus \{1\}$ has $\chi(G_{YO} \setminus \{1\}) = \chi_f(G_{YO} \setminus \{1\}) = \omega(G_{YO} \setminus \{1\}) = 3$ and yet its completion G_{YO_c} is a 25 vertex state-independent graph in dimension 3.

Conclusions. We have investigated the requirements for a set of projective measurements to reveal state-(in)dependent contextuality. In the state-dependent scenario, identifying a proper subgraph that is contextual in the realization is sufficient to guarantee contextuality of the graph. The condition for non-contextuality based on equality of $\theta(G)$ and $\alpha(G)$ does not hold for general orthogonality graphs.

By the rigorous measure of contextuality, the maximally mixed state was shown to be the least and pure states the most contextual among all states of a given Hilbert space dimension. A necessary and sufficient condition for state-independent contextuality was formulated in terms of the fractional chromatic number $\chi_f(G)$. A similarly formulated condition in terms of $\chi(G)$ fails to guarantee sufficiency of the S-IC property for G. $\chi_f(G)$ being in general easier to compute than $\chi(G)$, the formulated condition enables the identification of state-independent contextual measurements which have been utilized as resources in device-independent security [9]. Additional projectors required to realize complete measurements may form a cloud of contextuality around a graph G and enable it to possess state-independent properties. An interesting question emerges as to what conditions one may place on a graph G which by itself is not S-IC such that some of its optimally completed versions are.

Acknowledgements. We thank Prof. Ryszard Horodecki for a critical reading of the manuscript. This work is supported by ERC grant QOLAPS and the National Research Foundation & Ministry of Education, Singapore.

---

[*] Electronic address: ravishankar˙r@nus.edu.sg

[1] S. Kochen and E.P. Specker, J. Math. Mech. **17**, 59 (1967).

[2] A. A. Klyachko, M. A. Can, S. Binicoglu and A. S. Shumovsky, Phys. Rev. Lett. **101**, 020403 (2008).

[3] A. Cabello, L. E. Danielsen, A. J. Lopez-Tarrida and J. R. Portillo, arXiv/quant-ph:1211.5825 (2012).

[4] A. Cabello, arXiv/quant-ph:1112.5149 (2011).

[5] A. Cabello, J. M. Estebaranz and G. Garcia-Alcaine, Phys. Lett. A **339** 425 (2005).

[6] A. Peres, *Quantum Theory: Concepts and Methods* (Dordrecht: Kluwer), p 188, (1993).

[7] A. Cabello, Phys. Rev. Lett. **101** 210401 (2008).

[8] S. Yu and C. H. Oh, Phys. Rev. Lett. **108** 030402 (2012).

[9] K. Horodecki, M. Horodecki, P. Horodecki, R. Horodecki, M. Pawlowski and M. Bourennane, arXiv/quant-ph:1006.0468.

[10] T. Fritz, A. Leverrier and A. B. Sainz, arXiv/quant-ph:1212.4084 (2012).

[11] A. Grudka, K. Horodecki, M. Horodecki, P. Horodecki, R. Horodecki, P. Joshi, W. Klobus and A. Wojcik, arXiv/quant-ph:1209.3745 (2012).

[12] A. Cabello, S. Severini and A. Winter, arXiv/quant-ph:1010.2163 (2010).

[13] R. Ramanathan, A. Soeda, P. Kurzynski and D. Kaszlikowski, Phys. Rev. Lett. **109** 050404 (2012).

[14] C. Berge, Math.-Natur. Reihe **10** 114 (1961).

[15] M. Chudnovsky, N. Robertson, P. Seymour and R. Thomas, Annals of Mathematics **164** 1 (2006).

[16] L. Lovasz, IEEE Trans. Inf. Theory **25**, 1 (1979).

[17] P. Badziag, I. Bengtsson, A. Cabello, H. Granstrom and J. Larsson, Found. Phys. **41** (2011).

[18] P. J. Cameron, A. Montanaro, M. W. Newman, S. Severini and A. Winter, Electronic Journal of Combinatorics **14**, 1 (2007).

[19] E. R. Scheinerman and D. H. Ullman, *Fractional Graph Theory: A Rational Approach to the Theory of Graphs*, (Dover Publications, 2011).

[20] R. A. Horn and C. R. Johnson, *Matrix Analysis*, (Cambridge University Press, 1987).

Supplementary Material

Here, we present proofs of the propositions formulated in the main text.

I. Conditions for non-contextuality. To recognize whether perfect graphs are the only non-contextual graphs, necessary and sufficient conditions for non-contextuality need to be formulated. One possible condition is that $\theta(G) = \alpha(G)$ which rules out the violation of facet-defining inequalities of the kind $\sum_i \langle \Pi_i \rangle \leq \alpha(G)$. However, for simple non-weighted orthogonality graphs the example shown in Fig. (1) shows that this condition is not sufficient to ensure non-contextuality. For this graph, $\theta(G) = \alpha(G) = 3$, yet the set of projectors and state displayed in the figure lead to a maximum violation of one of the facet defining inequalities of G upto its maximum value of $\sqrt{5}$ showing that G is indeed (state-dependent) contextual. Furthermore, a numerical check via a semi-definite program reveals that the 25 vertex graph G_{YO_c} obtained by the unique completion of G_{YO} (Fig. (2)) considered in [8] has $\theta(G_{YOc}) = \alpha(G_{YOc}) = 13$ showing that the condition is not sufficient in the completion paradigm explained in the text as well.

Observation 1. If in any realization of orthogonal graph G, some proper subgraph H of G is contextual, then G is contextual.

Proof. Assume that for some proper subgraph H of an orthogonal graph G, there exists state ρ such that $\vec{x}^{\{\rho\}}(H) \notin STAB(H)$. Then $\vec{x}^{\{\rho\}}(G) \in STAB(G)$ implies that there exists a convex decomposition with probability distribution $\{p_j\}$ of the form

$$\vec{x}^{\{\rho\}}(G) = \sum_{j=1}^{|\mathcal{I}(G)|} p_j \vec{I}^j.$$

Assigning weights p_j to all independent sets $I^j(H) \subset I^j(G)$ with the independent set taken to be the null set for those $I^j(G)$ that do not contain any vertices of H, one obtains the non-contextual convex decomposition for $\vec{x}^{\{\rho\}}(H)$ which is a contradiction. Therefore, for any proper subgraph H of an orthogonal graph G, if there exists a state ρ such that $\vec{x}^{\{\rho\}}(H) \notin STAB(H)$, then $\vec{x}^{\{\rho\}}(G) \notin STAB(G)$. $\square$

Observation 2. Non-perfect graphs G containing induced C_5 are contextual in dimension 3.

Proof. In dimension 3, the most general class of projectors $\Pi_i = |v_i\rangle\langle v_i|$ realizing C_5 can be parametrized as [17] (in cyclic order): $|v_1\rangle = \{1,0,0\}^T$, $|v_2\rangle = \{0,1,0\}^T$, $|v_3\rangle = \{\cos\theta_1, 0, \sin\theta_1 e^{i\phi_1}\}^T$, $|v_4\rangle = \mathcal{N}\{e^{-i\phi_1}\sin\theta_1\cos\theta_2, e^{-i\phi_2}\sin\theta_2\cos\theta_1, -\cos\theta_1\cos\theta_2\}^T$, $|v_5\rangle = \{0, \cos\theta_2, \sin\theta_2 e^{i\phi_2}\}^T$. A numerical search then reveals that no combination of $\{\theta_1, \theta_2, \phi_1, \phi_2\}$ exists for which $\| \sum_{i=1}^5 |v_i\rangle\langle v_i| \|_\infty \leq 2$ if we require that the five projectors be distinct. In fact, as shown in [17] the number of parameters can be reduced to one using the fact that $\theta_1 = \theta_2 (= \theta, say)$ and $\phi_1 = \phi_2 = 0$ gives rise to the same spectrum for $\sum_{i=1}^5 |v_i\rangle\langle v_i|$. The infinity norm of this matrix is then given by

$$\kappa_1 = \frac{3+\sin^2\theta}{2} + \frac{1}{2}\sqrt{\frac{\sin^6\theta - 5\sin^4\theta + 3\sin^2\theta + 1}{1+\sin^2\theta}}.$$

The above expression only decreases to 2 for $\sin\theta = \{0, 1\}$ in which case the projectors are no longer distinct. The fact that the only facet-defining inequality of $STAB(C_5)$ that is not also facet-defining for $QSTAB(C_5)$ is given by $\sum_{i=1}^5 \langle |v_i\rangle\langle v_i| \rangle \leq \alpha(C_5) = 2$ then proves that when C_5 is realized in dimension 3, irrespective of the particular set of projectors used, there always exists a quantum state whose corresponding vector of expectation values lies outside the non-contextual polytope $STAB(C_5)$. Therefore by Observation 1, G containing induced C_5 is contextual for dimension 3. $\square$

Theorem 1. $M^{(G)}(\rho)$ is entirely determined by the spectrum $\vec{\kappa}(\rho)$. Moreover, $\forall \rho_1, \rho_2 \in \mathcal{H}^d$ and $\forall G$, $\vec{\kappa}(\rho_1) \prec \vec{\kappa}(\rho_2)$ implies $M^{(G)}(\rho_1) \leq M^{(G)}(\rho_2)$. In particular, $M^{(G)}(\mathbb{1}_d) \leq M^{(G)}(\rho)\ \forall \rho \in \mathcal{H}^d$.

Proof. The fact that the measure of contextuality $M^{(G)}(\rho)$ is determined by $\vec{\kappa}(\rho)$ is a consequence of the maximization procedure in Eq.(2). Two states η_1, η_2 sharing the same spectrum $\vec{\kappa}(\eta)$ are unitarily equivalent, $\eta_1 = U\eta_2 U^\dagger$ for some unitary U. Therefore, given a set of projectors $\{\Pi_k\}(\eta_1)$ that result from the maximization in the computation of the measure for η_1, one may construct a corresponding set of projectors $\{U^\dagger \Pi_k U\}$ that gives the same value of the measure for η_2 (and vice versa). The sole dependence of $M^{(G)}(\rho)$ on the spectrum $\vec{\kappa}(\rho)$ follows.

Given $\rho_1, \rho_2 \in \mathcal{H}^d$, if their vector of eigenvalues obey the majorization relation $\vec{\kappa}(\rho_1) \prec \vec{\kappa}(\rho_2)$, it then follows



FIG. 1: An example graph for which $\theta(G) = \alpha(G) = 3$ which is nevertheless a state-dependent contextual graph. For instance, consider the projectors corresponding to $|v_1\rangle = \{0,1,0,0\}^T$, $|v_2\rangle = \{0,0,\cos\theta,\sin\theta\}^T$, $|v_3\rangle = \mathcal{N}\{0,\cos\theta\sin\theta,\cos\theta\sin\theta,-\cos^2\theta\}^T$, $|v_4\rangle = \{0,\cos\theta,0,\sin\theta\}^T$, $|v_5\rangle = \{0,0,1,0\}^T$, $|v_6\rangle = \{\cos\phi,\sin\phi,0,0\}^T$ with $\sin\theta = \sqrt{(1/2)(\sqrt{5}-1)}$ and $\phi = \pi/4$ which lead to a faithful realization of the graph. Then the state $|\psi\rangle = \{0,\frac{1}{5^{1/4}},\frac{1}{5^{1/4}},\frac{1}{\sqrt{5+2\sqrt{5}}}\}^T$ with these projectors gives a violation of the facet of STAB(G) defined by the inequality $\sum_{i=1}^{5}\langle|v_i\rangle\langle v_i|\rangle \leq 2$ upto its maximum value of $\sqrt{5}$.

[20] that there exist d-dimensional permutation matrices P_j and a probability distribution $\{p_j\}$ such that

$$\vec{\kappa}(\rho_1) = \sum_j p_j P_j \vec{\kappa}(\rho_2).$$

Given an orthogonal graph G, let us denote the set of corresponding projectors resulting from the maximization in the measure $M^{(G)}(\rho_1)$ by $\{\Pi_k\}(\rho_1)$ as above. With ρ_2 written in diagonal form, let $\rho_2^j = P_j\rho_2 P_j^\dagger$ denote the diagonal matrices obtained from the permutation of the eigenbasis of ρ_2. For each ρ_2^j, let the joint probability distribution resulting from the minimization in Eq.(2) with the chosen set of projectors $\{\Pi_k\}(\rho_1)$ be denoted by $p^{\rho_2^j}(\lambda)$. We then have,

$$\begin{aligned}
M^{(G)}(\rho_2) \geq \sum_j p_j M^{(G)}_{\{\Pi_k\}(\rho_1)}(\rho_2^j) &= \sum_{c\in\mathcal{C}(G)} \frac{1}{|\mathcal{C}(G)|} \sum_j p_j \left(D(q^{(\rho_2^j,\{\Pi_k\}(\rho_1))}(\lambda_c) \| p^{\rho_2^j}(\lambda_c)) \right) \\
&\geq \sum_{c\in\mathcal{C}(G)} \frac{1}{|\mathcal{C}(G)|} \left(D(\sum_j p_j q^{(\rho_2^j,\{\Pi_k\}(\rho_1))}(\lambda_c) \| \sum_j p_j p^{\rho_2^j}(\lambda_c)) \right) \\
&\geq \sum_{c\in\mathcal{C}(G)} \frac{1}{|\mathcal{C}(G)|} \left(D(q^{(\rho_1,\{\Pi_k\}(\rho_1))|}(\lambda_c) \| \sum_j p_j p^{\rho_2^j}(\lambda_c)) \right) \\
&\geq M^{(G)}(\rho_1).
\end{aligned}$$

In the above, the first inequality is a result of the fact that $\{\Pi_k\}(\rho_1)$ may not yield the maximum in the computation of the measure for ρ_2, the second inequality is a result of the convexity property of the relative entropy, the third follows from linearity, and the fourth by the definition of the measure for state ρ_1. The monotonicity of the measure of contextuality with majorization order thus follows. Since the spectrum of the maximally mixed state $\mathbb{1}_d$ is majorized by that of every other state $\rho \in \mathcal{H}^d$, it follows that $\mathbb{1}_d$ is the least contextual among all states belonging to a given Hilbert space and analogously, pure states are the most contextual. □

Proposition 1. *C_{2k+1} and $\bar{C}_{2k+1}$ do not represent state-independent contextual measurements for any integral k.*

Proof. That $\vec{x}^{\{\mathbb{1}_d\}}(C_{2k+1}) \in STAB(C_{2k+1})$ can be easily seen by noting that $\chi(C_{2k+1}) = 3$ for all integral k. The three colour classes in a proper coloring of C_{2k+1} thus form a partition of the vertex set of C_{2k+1} into 3 independent sets I_1, I_2 and I_3. For any dimension $d \geq 3$ in which the graphs are realized by rank-1 projectors therefore, the vector $\vec{x}^{\{\mathbb{1}_d\}}(C_{2k+1})$ (which is a uniform vector with $\vec{x}_j^{\{\mathbb{1}_d\}}(C_{2k+1}) = \frac{1}{d}$ for all $1 \leq j \leq 2k+1$) can be decomposed as a convex combination with weights $1/d$ of the incidence vectors for the three independent sets $\vec{I}^{(1)}(C_{2k+1}), \vec{I}^{(2)}(C_{2k+1}), \vec{I}^{(3)}(C_{2k+1})$ and weight $(1-3/d)$ for the incidence vector $\vec{I}^{(0)}(C_{2k+1})$ with 0's for all $2k+1$ vertices, i.e.,

$$\vec{x}^{\{\mathbb{1}_d\}}(C_{2k+1}) = \frac{1}{d}\left(\vec{I}^{(1)} + \vec{I}^{(2)} + \vec{I}^{(3)}\right) + (1-\frac{3}{d})\vec{I}^{(0)}.$$

For the complements of odd cycles, $\chi(\bar{C}_{2k+1}) = k+1$ while $\omega(\bar{C}_{2k+1}) = k$. However, these graphs cannot be realized by distinct rank-1 projectors in dimension k which can be seen as follows. Label the vertices of C_{2k+1} from 1 to $2k+1$ in cyclic order. In the complementary graph $\bar{C}_{2k+1}$, the vertex set $\{1, 3, \ldots, 2k-3, 2k-1\}$ forms a clique of size k as does the vertex set $\{1, 3, \ldots, 2k-3, 2k\}$. In order for this graph to be realized in dimension k therefore, vertices labeled $2k-1$ and $2k$ must be assigned the same projector orthogonal to the $k-1$ dimensional space spanned by the rest of the vertices $\{1, \ldots, 2k-3\}$. This contradicts the requirement that distinct vertices be assigned distinct projectors, therefore the minimal dimension in which $\bar{C}_{2k+1}$ can be realized is at least $k+1$. From the previous construction, $\vec{x}^{\{\mathbb{1}_d\}}(\bar{C}_{2k+1})$ admits a convex decomposition into non-contextual assignments (which are incidence vectors of independent sets in a proper coloring) for any $d \geq k+1$ and therefore, these graphs cannot be state-independent for any k as well. □

Theorem 2. An orthogonal graph G realizable by a set of rank-r projective measurements $\{\Pi_j\}$ with $\Pi_j \in \mathcal{H}^d$ is state-independent contextual for dimension d if and only if $\chi_f(G) > \frac{d}{r}$.

Proof. For given set of rank-r projective measurements $\{\Pi_j\}$ in d-dimensional space, let us construct the orthogonality graph G with a total of $|\{\Pi_j\}| = n$ vertices. The vector of expectation values for the maximally mixed state $\vec{x}^{\{\mathbb{1}_d\}}(G)$ is then the uniform vector with components $\vec{x}_k^{\{\mathbb{1}_d\}}(G) = \frac{r}{d}$ for $1 \leq k \leq n$. The fractional chromatic number is by definition $\chi_f(G) = \min \frac{a}{b}$ where out of a total of $'a'$ colors, $'b'$ colors are assigned to each vertex such that vertices connected by an edge receive disjoint sets of colors. Let $\{a^*(G), b^*(G)\}$ denote the values of a and b that achieve the minimum in this definition. For each of the colors $1 \leq c \leq a^*(G)$, let $\vec{I}^{(c)}(G)$ denote the incidence vector of the independent set of vertices colored with color c and $\vec{I}^{(0)}(G)$ denote the vector with all components 0. We may consider three cases.

(i) $\chi_f(G) = \frac{a^*}{b^*} = \frac{d}{r}$.

In this case, the fractional colorings themselves yield the convex decomposition

$$\vec{x}^{\{\mathbb{1}_d\}}(G) = \sum_{c=1}^{a^*} \frac{1}{a^*}\vec{I}^{(c)}(G).$$

Since each vertex in the decomposition appears in a total of b^* independent sets, the uniform vector representing the maximally mixed state is exactly reproduced.

(ii) $\chi_f(G) = \frac{a^*}{b^*} < \frac{d}{r}$.

Let $a' = \frac{b^* d}{r} > a^*$. Then one may construct the following decomposition into non-contextual assignments

$$\vec{x}^{\{\mathbb{1}_d\}}(G) = \sum_{c=1}^{a^*} \frac{1}{a'}\vec{I}^{(c)}(G) + (1-\frac{a^*}{a'})\vec{I}^{(0)}(G)$$

with the final vector appearing to ensure the probabilities sum to unity. Therefore, once again we have that $\vec{x}^{\{\mathbb{1}_d\}}(G) \in STAB(G)$.

(iii) $\chi_f(G) > \frac{d}{r}$.

To prove that in this case, we definitely have $\vec{x}^{\{\mathbb{1}_d\}}(G) \notin STAB(G)$, we may use the formulation of $\chi_f(G)$ in terms of a linear program. Let us assign weights w_j to each independent set $I^{(j)}(G) \in \mathcal{I}(G)$. Then $\chi_f(G) = min\{\sum_{j=1}^{|\mathcal{I}(G)|} w_j : \sum_{j=1}^{|\mathcal{I}(G)|} w_j \vec{I}^{(j)}(G) \geq \vec{\mathbf{1}}\}$ where $\vec{\mathbf{1}}$ denotes the uniform vector of 1's on each vertex. The following Lemma [19] then enables us to change the inequality in the definition to equality.

Lemma: If a graph G has a fractional coloring with total weight $w = \sum_{j=1}^{|\mathcal{I}(G)|} w_j$ such that $\sum_{j=1}^{|\mathcal{I}(G)|} w_j \vec{I}^{(j)}(G) \geq \vec{\mathbf{1}}\}$, then one may construct a fractional coloring with weight $w' \leq w$ such that $\sum_{j=1}^{|\mathcal{I}(G)|} w'_j \vec{I}^{(j)}(G) = \vec{\mathbf{1}}\}$.

Assume $\vec{x}^{\{\mathbb{1}_d\}}(G) \in STAB(G)$, and let the following be a convex decomposition for $\vec{x}^{\{\mathbb{1}_d\}}(G)$ into non-contextual assignments.

$$\vec{x}^{\{\mathbb{1}_d\}}(G) = \sum_{j=1}^{\mathcal{I}(G)} p_j \vec{I}^{(j)}(G).$$

Then $\sum_{j=1}^{\mathcal{I}(G)} \frac{p_j d}{r}\vec{I}^{(j)} = \vec{\mathbf{1}}$ so that $\chi_f(G) \leq \sum_{j=1}^{\mathcal{I}(G)} \frac{p_j d}{r} = \frac{d}{r}$ which is a contradiction. Therefore, $\chi_f(G) > \frac{d}{r}$ is a necessary and sufficient condition for $\vec{x}^{\{\mathbb{1}_d\}}(G) \notin STAB(G)$ (in other words, for G to be state-independent contextual). □

II. An explicit decomposition of $\vec{x}^{\{\mathbb{1}_7\}}(J(G_{YO}, G_{YO}))$. Consider two copies of the state-independent graph G_{YO} considered in [8] and reproduced in Fig. (2). The join of the two copies is defined as the graph $J(G_{YO}, G_{YO})$ obtained by the union of the two graphs with additional edges connecting every vertex of the first copy with every vertex of the second.

Let us label the vertices $\{1, \ldots, 13\}$ in one copy as in Fig. (2) and $\{1', \ldots, 13'\}$ in the second copy. For notational convenience, let us use $\vec{I}(\{v_i\})$ to denote the incidence vector of the independent set $\{v_i\}$. $J(G_{YO}, G_{YO})$ has $\omega(J(G_{YO}, G_{YO})) = 6$ and $\chi(J(G_{YO}, G_{YO})) = 8$ so one may expect it to state-independent contextual (by itself without completion) for dimension 7. However,



FIG. 2: The 13 vertex graph G_{YO}. The join of two copies of this graph provides a counter-example to $\chi(G) > d$ being a sufficient condition for a graph to be state-independent contextual by itself.

the following explicit convex decomposition of the vector $\vec{x}^{\{\mathbb{1}_7\}}(J(G_{YO}, G_{YO}))$ proves otherwise.

$$\begin{aligned}\vec{x}^{\{\mathbb{1}_7\}}(J(G_{YO}, G_{YO})) \;=\; & \frac{1}{77}\big[5\vec{I}(\{1,5,6,10\}) + 4\vec{I}(\{1,5,9,12\}) + \vec{I}(\{1,6,8,11\}) + \vec{I}(\{1,10,11,12,13\}) + 2\vec{I}(\{2,4,6,11\}) \\ + \;& 2\vec{I}(\{2,4,9,12\}) + 3\vec{I}(\{2,6,7,10\}) + 3\vec{I}(\{2,7,9,13\}) + \vec{I}(\{2,10,11,12,13\}) + 5\vec{I}(\{3,4,8,11\}) \\ + \;& 5\vec{I}(\{3,7,8,13\}) + \vec{I}(\{3,10,11,12,13\}) + 2\vec{I}(\{4,5,9,12\})\big] + \frac{1}{77}\,(v_i \leftrightarrow v_i') + \frac{1}{11}\vec{I}^0\end{aligned}$$

Here, $(v_i \leftrightarrow v_i')$ denotes the same combination of independent sets considered on the second copy of G_{YO} and $\vec{I}^0$ denotes the vector with all vertices assigned value 0. From the above, note that $\chi_f(J(G_{YO}, G_{YO})) = 6\frac{4}{11}$. Note that this does not rule out the possibility that a completed version of $J(G_{YO}, G_{YO})$ in dimension 7 is a state-independent graph.

Proposition 2. An orthogonal graph G can be state-independent contextual by itself only for dimensions $\xi(G) \leq d < 2^{\xi(G)-1}$.

Proof. By definition, $\xi(G) = d$ denotes the smallest dimension in which the orthogonal graph G on n vertices is realized by rank-1 projectors with distinct projectors assigned to distinct vertices and two vertices connected by an edge if and only if the corresponding projectors are orthogonal. The proof proceeds by showing that for all G, $\chi(G) \leq 2^{\xi(G)-1}$ so that the result follows from the necessity (not sufficiency) of $\chi(G) > d$ for G to be state-independent contextual. Let $\{\Pi_k = |v_k\rangle\langle v_k|\} \subset \mathcal{H}^d$ denote a particular realization of G with the components of the column vector $|v_k\rangle$ being in general given by $|v_k\rangle^{(j)} = |v_k^{(j)}|e^{i\theta_k^{(j)}}$ for $1 \leq j \leq (d-1)$ and $|v_k\rangle^{(d)} = |v_k^{(d)}|$. Firstly note that by a suitable rotation of $\{\Pi_k\}$, it is possible to ensure that $\forall k$, the component $|v_k\rangle^{(d)} \neq 0$. Let us describe each of the components $|v_k\rangle^{(j)}$ by the quadrant in which $\theta_k^{(j)}$ lies: each $|v_k\rangle^{(j)}$ is then represented by a 4-dimensional dit taking value $t_k^{(j)}$ $(1 \leq t_k^{(j)} \leq 4)$ according to the quadrant $(t_k^{(j)}-1)\frac{\pi}{2} \leq \theta_k^{(j)} < t_k^{(j)}\frac{\pi}{2}$. Each $|v_k\rangle$ then belongs to one of the 4^{d-1} quadrant classes created by the value of the corresponding vector $\vec{t}_k = \{t_k^{(1)}, \ldots, t_k^{(d)}\}^T$. Since no two $|v_k\rangle$ belonging to the same quadrant class can be orthogonal, this results in a partition of the vertex set into 4^{d-1} independent sets each of which can be assigned its own color in a proper coloring. Finally, noting that no two $|v_k\rangle$ with all entries restricted to the first and fourth quadrants, i.e., all $t_k^{(j)} \in \{1,4\}$ (or equivalently to the second and third) can be orthogonal, one may partition the vertex set into a total of 2^{d-1} independent sets. Within each independent set, all vertices may be assigned the same color so that a proper coloring of G requires at most $2^{\xi(G)-1}$ colors. By the construction of the non-contextual decomposition of $\vec{x}^{\{\mathbb{1}_d\}}(G)$ with the incidence vectors of the independent sets in a proper coloring (together with the vector of all zeros $\vec{I}^{(0)}(G)$), we see that G cannot be an S-IC graph for $d \geq 2^{\xi(G)-1}$ □

Proposition 3. All basis-complete graphs G_c for which $\chi_f(G_c) = \omega(G_c)$ $(= n$, say) are proper subgraphs of the Kneser graphs $K(nb : b)$ for some integral b. $K(nb : b)$ with $n \geq 2$ and $b \geq 2$ themselves cannot be quantumly realized by rank-1 projectors in dimension n. Moreover for $b = 2$, all proper basis-complete subgraphs of $K(2n : 2)$ also cannot be realized in dimension n.

Proof. Firstly, notice that a basis-complete graph G_c that has $\chi_f(G_c) = \frac{a}{b} = \omega(G_c)$ must have a fractional coloring with all a colors used in each clique and b colors assigned to each vertex of the clique. For $\omega(G_c) = n$ therefore, they are precisely the subgraphs of $K(nb : b)$ by definition since the Kneser graphs contain all possible b element subsets of the $a = nb$ element set.

In dimension $n = 2$, no graph apart from a single edge is quantumly realizable by rank-1 projectors. Assume that a Kneser graph $K(nb : b)$ with $n \geq 3$ for some $b \geq 2$ is realizable in dimension n by distinct rank-1 projectors. Each vertex is assigned b colors out of the vertex set $\{1, \ldots, nb\}$ and all b element sets are considered, so we may without loss of generality assign projectors to b element color sets. Moreover, we may assign to the color set $S_k = \{1+(k-1)b, \ldots, kb\}$ for $1 \leq k \leq n$ the projector Π_k onto dimension k, so that the set $\{S_k\}$ is represented by the standard computational basis in dimension n. Let us now partition one of the color sets, S_1 say, into two non-empty subsets $S_1^{(2)}$ and $S_1^{(3)}$ of size b_1 and b_2 respectively, $|S_1^{(2)}| = b_1$ and $|S_1^{(3)}| = b_2$. A new b-element set $S_{1,2}$ can be formed by adding $b - b_1$ colors from S_2 to $S_1^{(2)}$ and similarly $S_{1,3}$ can be formed by adding $b - b_2$ colors from S_3 to $S_1^{(3)}$. These new sets $S_{1,2}$ and $S_{1,3}$ must also by definition be represented by vertices in $K(nb : b)$. Now, by orthogonality, the projector $\Pi_{1,2}$ assigned to $S_{1,2}$ belongs to the subspace formed by the span of Π_1 and Π_2 and similarly the projector $\Pi_{1,3} \in span\{\Pi_1, \Pi_3\}$. However, by construction $\Pi_{1,2}$ and $\Pi_{1,3}$ are orthogonal so that either $\Pi_{1,2} = \Pi_2$ or $\Pi_{1,3} = \Pi_3$ which provides a contradiction to the requirement of distinct projectors for distinct vertices.

Finally, let us show that basis-complete subgraphs G_c of $K(2n : 2)$ also cannot be realized in dimension n. Assume that a subgraph G_c exists that can be realized in dimension n. The following statements can be made without loss of generality. As before, consider the clique $C_1 = \{v_k\}$ in G_c with color set $\{S_k\}$ $(S_k = \{2k-1, 2k\})$ for $1 \leq k \leq n$ and assign the computational basis projectors Π_k onto dimension k to its vertices. Note that every clique must contain all the colors $\{1, \ldots, 2n\}$. Now, consider a second clique C_2 containing two vertices v'_2 and v'_3 not belonging to C_1. Without loss of generality, let the colors assigned to v'_2 be $\{1, 3\}$ and the color 2 be assigned to v'_3. If the color set of v'_3 is $S'_3 = \{2, 4\}$, then the corresponding projectors $\Pi'_2 \perp \Pi'_3$ that belong to span of Π_1 and Π_2 may be realized. However, if every such pair of vertices has similar assignments, then $\chi(G_c) = \omega(G_c)$. Therefore, there must exist vertex $v'_3 \in C_2$ which is assigned color set $S'_3 = \{2, c\}$ with $c \notin \{1, 2, 4\}$, $c = 5$ say. Then, the corresponding projectors $\Pi'_2 \in \text{span}\{\Pi_1, \Pi_2\}$ and $\Pi'_3 \in \text{span}\ \{\Pi_1, \Pi_3\}$ which results either $\Pi'_2 = \Pi_2$ or $\Pi'_3 = \Pi_3$. Again, the requirement of distinct projectors in $\mathcal{H}^n$ being assigned to distinct vertices of $K(2n : 2)$ cannot be met for all n. □